

A NONCOMMUTATIVE SYMMETRIC SYSTEM OVER THE GROSSMAN-LARSON HOPF ALGEBRA OF LABELED ROOTED TREES

WENHUA ZHAO

ABSTRACT. In this paper, we construct explicitly a noncommutative symmetric (NCS) system over the Grossman-Larson Hopf algebra of labeled rooted trees. By the universal property of the NCS system formed by the generating functions of certain noncommutative symmetric functions, we obtain a specialization of noncommutative symmetric functions by labeled rooted trees. Taking the graded duals, we also get a graded Hopf algebra homomorphism from the Connes-Kreimer Hopf algebra of labeled rooted forests to the Hopf algebra of quasi-symmetric functions. A connection of the coefficients of the third generating function of the constructed NCS system with the order polynomials of rooted trees is also given and proved.

1. Introduction

Let K be any unital commutative $\mathbb{Q}$ -algebra and A a unital associative but not necessarily commutative K -algebra. Let t be a formal central parameter, i.e. it commutes with all elements of A , and $A[[t]]$ the K -algebra of formal power series in t with coefficients in A . A *NCS* (*noncommutative symmetric*) *system* over A (see Definition 2.1) by definition is a 5-tuple $\Omega \in A[[t]]^{\times 5}$ which satisfies the defining equations (see Eqs. (2.1)–(2.5)) of the NCSFs (noncommutative symmetric functions) first introduced and studied in the seminal paper [GKLLRT]. When the base algebra K is clear in the context, the ordered pair (A, Ω) is also called a *NCS system*. In some sense, a NCS system over an associative K -algebra can be viewed as a system of analogs in A of the NCSFs defined by Eqs. (2.1)–(2.5). For some general discussions on

Date: February 2, 2008.

2000 Mathematics Subject Classification. Primary: 05E05, 16W30; Secondary: 06A07, 06A11 .

Key words and phrases. NCS systems, noncommutative symmetric functions, quasi-symmetric functions, specializations, the Grossman-Larson Hopf algebra, the Connes-Kreimer Hopf algebra, labeled rooted trees, the (strict) order polynomials of posets.

the NCS systems, see [Z2]. For a family of NCS systems over differential operator algebras and their applications to the inversion problem, see [Z3] and [Z4]. For more studies on NCSFs, see [T], [KLT], [DKKT], [KT1], [KT2] and [DFT].

One immediate but probably the most important example of the NCS systems is $(NSym, \Pi)$ formed by the generating functions of the NCSFs defined in [GKLLRT] by Eqs. (2.1)–(2.5) over the free K -algebra $NSym$ of NCSFs (see Section 2). It serves as the universal NCS system over all associative K -algebra (see Theorem 2.6). More precisely, for any NCS system (A, Ω) , there exists a unique K -algebra homomorphism $\mathcal{S} : NSym \rightarrow A$ such that $\mathcal{S}^{\times 5}(\Pi) = \Omega$ (here we have extended the homomorphism $\mathcal{S}$ to $\mathcal{S} : NSym[[t]] \rightarrow A[[t]]$ by the base extension).

The universal property of the NCS system $(NSym, \Pi)$ can be applied as follows when a NCS system (A, Ω) is given. Note that, as an important topic in the symmetric function theory, the relations or polynomial identities among various NCSFs have been worked out explicitly (see [GKLLRT]). When we apply the K -algebra homomorphism $\mathcal{S} : NSym \rightarrow A$ guaranteed by the universal property of the system $(NSym, \Pi)$ to these identities, they are transformed into identities among the corresponding elements of A in the system Ω . This will be a very effective way to obtain identities for certain elements of A if we can show they are involved in a NCS system over A . On the other hand, if a NCS system (A, Ω) has already been well-understood, the K -algebra homomorphism $\mathcal{S} : NSym \rightarrow A$ in return provides a *specialization* or *realization* ([GKLLRT], [St2]) of NCSFs, which may provide some new understandings on NCSFs. For more studies on the specializations of NCSFs, see the references quoted above for NCSFs.

In this paper, we apply the gadget above to the Grossman-Larson Hopf algebra of labeled rooted trees. To be more precise, for any non-empty $W \subseteq \mathbb{N}^+$,¹ let $\mathcal{H}_{GL}^W$ be the Grossman-Larson Hopf algebra $\mathcal{H}_{GL}^W$ ([GL], [CK], [F]) of rooted trees labeled by positive integers of W . We first introduce five generating functions (see Eqs. (4.1)–(4.4) and Eq. (4.15)) of certain elements of $\mathcal{H}_{GL}^W$ and show that they form a NCS system $\Omega_{\mathbb{T}}^W$ over $\mathcal{H}_{GL}^W$ (see Theorem 4.5). Then, by the universal property of the NCS system $(NSym, \Pi)$ from NCSFs, we obtain a graded Hopf algebra homomorphism $\mathcal{T}_W : NSym \rightarrow \mathcal{H}_{GL}^W$, which gives a specialization of NCSFs by W -labeled rooted trees (see Theorem 4.6).

¹All constructions and results of this paper work equally well for any non-empty weighted set W , i.e. any non-empty set W with a fixed weight function $wt : W \rightarrow \mathbb{N}^+$ such that, for any $k \in \mathbb{N}^+$, $wt^{-1}(k)$ is a finite subset of W . But, for simplicity and convenience, we will always assume that W is a non-empty subset of $\mathbb{N}^+$.

By taking the graded duals, we get a graded Hopf algebra homomorphism $\mathcal{T}_W^* : \mathcal{H}_{CK}^W \rightarrow \mathcal{QSym}$ from the Connes-Kreimer Hopf algebra $\mathcal{H}_{CK}^W$ ([Kr], [CK], [F]) to the Hopf algebra $\mathcal{QSym}$ of quasi-symmetric functions ([Ge], [MR], [St2]). Later in [Z5], it will be shown that, when $W = \mathbb{N}^+$, the specialization $\mathcal{T}_W$ above is actually an embedding and hence the Hopf algebra homomorphism $\mathcal{T}_W^*$ is onto. Finally, we give a combinatorial interpretation of the constants θ_T (see Definition 4.2) of rooted trees T that appeared in the third component $\tilde{d}(t)$ (see Eq. (4.15)) of the NCS system $\Omega_{\mathbb{T}}^W$ above. We show that, for each rooted tree T , the constant θ_T coincides with the coefficient of s in the order polynomial $\Omega(B_-(T), s)$ (see [St2]), where $B_-(T)$ is the rooted forest obtained by cutting off the root of T .

The arrangement of the paper is as follows. In Section 2, we first recall the definition of the NCS systems ([Z2]) Ω over the K -algebras A and a result (see Proposition 2.2) on the NCS systems when A is further a bialgebra or Hopf algebra. We then recall the universal NCS system $(\mathcal{NSym}, \Pi)$ formed by generating functions of certain NCSFs in [GKLLRT]. The Hopf algebra structure of $\mathcal{NSym}$ and the universal property of the NCS system $(\mathcal{NSym}, \Pi)$ (see Theorem 2.6) will also be reviewed.

In Section 3, we first fix certain notation on rooted trees and recall the Connes-Kreimer Hopf algebra $\mathcal{H}_{CK}^W$ and the Grossman-Larson Hopf algebra $\mathcal{H}_{GL}^W$ of W -labeled rooted forests and rooted trees, respectively. Then, by using the duality between the Grossman-Larson Hopf algebra and the Connes-Kreimer Hopf algebra (see Theorem 3.2), we prove a technic lemma, Lemma 3.4, that will be crucial for our later arguments.

In Section 4, we introduce five generating functions of W -labeled rooted trees and show in Theorem 4.5 that they form a NCS system $\Omega_{\mathbb{T}}^W$ over the Grossman-Larson Hopf algebra $\mathcal{H}_{GL}^W$. By the universal property of the system $(\mathcal{NSym}, \Pi)$, we get a graded K -Hopf algebra homomorphism $\mathcal{T}_W : \mathcal{NSym} \rightarrow \mathcal{H}_{GL}^W$ (see Theorem 4.6). By taking the graded duals, we get a graded Hopf algebra homomorphism $\mathcal{T}_W^* : \mathcal{H}_{CK}^W \rightarrow \mathcal{QSym}$ from the Connes-Kreimer Hopf algebra $\mathcal{H}_{CK}^W$ to the Hopf algebra $\mathcal{QSym}$ of quasi-symmetric functions (see Corollary 4.7).

In Section 5, we first recall the strict order polynomials and the order polynomials of finite posets (partially ordered sets). Then, by applying some of results proved in [WZ] for the strict order polynomials of rooted forests and the well-known Reciprocity Relation (see Proposition 5.1) between the strict order polynomials and the order polynomials of finite posets, we show in Proposition 5.8 that, for any $T \in \mathbb{T}$, the constant θ_T involved in the third component of the NCS system $\Omega_{\mathbb{T}}^W$ is same as

the coefficient of s of the order polynomial $\Omega(B_-(T), s)$ of the rooted forest $B_-(T)$.

Finally, two remarks are as follows. First, as we pointed out early, by applying the specialization $\mathcal{T}_W : \mathcal{NSym} \rightarrow \mathcal{H}_{GL}^W$, we will get a host of identities for the rooted trees involved in the NCS system $\Omega_{\mathbb{T}}^W$ from the identities of the NCSFs in II. We believe some of these identities are interesting, at least from a combinatorial point view. But, to keep this paper in a certain size, we have to ask the reader who is interested to do the translations via the Hopf algebra homomorphism $\mathcal{T}_W : \mathcal{NSym} \rightarrow \mathcal{H}_{GL}^W$. Secondly, some relations between the NCS system $(\mathcal{H}_{GL}^W, \Omega_{\mathbb{T}}^W)$ constructed in this paper and the NCS systems constructed in [Z3] over differential operator algebras will be further studied in the followed paper [Z5]. Some consequences of those relations to the inversion problem ([BCW] and [E]) and specializations of NCSFs will also be derived there. In particular, it will be shown that, with the label set $W = \mathbb{N}^+$, the K -Hopf algebra homomorphism $\mathcal{T}_W : \mathcal{NSym} \rightarrow \mathcal{H}_{GL}^W$ in Theorem 4.6 is actually an embedding.

Acknowledgment: The author is very grateful to both referees for many invaluable suggestions to improve the paper. In particular, all the diagrams of trees in this paper are due to one of the referees who very kindly sent the author all the latex commands.

2. The Universal NCS System from Noncommutative Symmetric Functions

In this section, we first recall the definition of the NCS systems ([Z2]) over associative algebras and some of the NCSFs (noncommutative symmetric functions) first introduced and studied in the seminal paper [GKLLRT]. We then discuss the universal property of the NCS system formed by the generating functions of these NCSFs. The main result that we will need later is Theorem 2.6 which was proved in [Z2]. For some general discussions on the NCS systems, see [Z2]. For more studies on NCSFs, see [T], [KLT], [DKKT], [KT1], [KT2] and [DFT].

Let K be any unital commutative $\mathbb{Q}$ -algebra² and A any unital associative but not necessarily commutative K -algebra. Let t be a formal central parameter, i.e. it commutes with all elements of A , and $A[[t]]$ the K -algebra of formal power series in t with coefficients in A . First let us recall the following main notion of this paper.

²For the reader who is mainly interested in the combinatorial aspects of the main results of this paper, the base field K throughout this paper can be safely chosen to be the field $\mathbb{Q}$ of rational numbers.

Definition 2.1. ([Z2]) *For any unital associative K -algebra A , a 5-tuple $\Omega = (f(t), g(t), d(t), h(t), m(t)) \in A[[t]]^{\times 5}$ is said to be a NCS (noncommutative symmetric) system over A if the following equations are satisfied.*

$$(2.1) \quad f(0) = 1$$

$$(2.2) \quad f(-t)g(t) = g(t)f(-t) = 1,$$

$$(2.3) \quad e^{d(t)} = g(t),$$

$$(2.4) \quad \frac{dg(t)}{dt} = g(t)h(t),$$

$$(2.5) \quad \frac{dg(t)}{dt} = m(t)g(t).$$

When the base algebra K is clear in the context, we also call the ordered pair (A, Ω) a *NCS system*. Since NCS systems often come from generating functions of certain elements of A that are under the consideration, the components of Ω will also be refereed as the *generating functions* of their coefficients.

All K -algebras A that we are going to work on in this paper are K -Hopf algebras. We will freely use some standard notions and results from the theory of bialgebras and Hopf algebras, which can be found in the standard text books [A], [Knu] and [Mo]. For example, by a *sequence of divided powers* of a bialgebra or Hopf algebra A we mean a sequence $\{c_n \mid n \geq 0\}$ of elements of A such that, for any $n \geq 0$, we have

$$\Delta c_n = \sum_{k \geq 0} c_k \otimes c_{n-k}.$$

The following result proved in [Z2] later will be useful to us.

Proposition 2.2. *Let (A, Ω) be a NCS system as above. Suppose A is further a K -bialgebra. Then the following statements are equivalent.*

- (a) *The coefficients of $f(t)$ form a sequence of divided powers of A .*
- (b) *The coefficients of $g(t)$ form a sequence of divided powers of A .*
- (c) *One (hence also all) of $d(t)$, $h(t)$ and $m(t)$ has all its coefficients primitive in A .*

In the following remark, we would like to point out a connection of the notion of NCS systems with the notion of combinatorial Hopf algebras, which was first introduced by M. Aguiar, N. Bergeron and F. Sottile in [ABS].

Remark 2.3. *First, as pointed out in Remark 2.17 in [Z2], when A is a graded and connected Hopf algebra, and one of the statements of*

Proposition 2.2 holds, say statement (b). Furthermore assume in this case that the coefficients of t^m ($m \geq 0$) of $g(t)$ are homogeneous with grading m . Then the data $(A, g(t))$ is equivalent to a combinatorial Hopf algebra structure on the graded dual Hopf algebra A^* of A . For more details of the equivalence above, see Remark 2.17 in [Z2].

Since all other components of Ω are completely determined by $g(t)$ (see Lemma 2.5 of [Z2] or Theorem 2.6 below), the notion of NCS systems under the conditions above is also equivalent to the notion of combinatorial Hopf algebras. Therefore, from this point of view, the notion of NCS systems generalizes the notion of combinatorial Hopf algebras to associative K -algebras A , since, for NCS systems over A , A does not have to be a bialgebra nor Hopf algebra and the equivalent conditions in Proposition 2.2 do not have to hold either.

On the other hand, we would also like to point out that the notion of NCS emphasizes the whole package of five generating functions of elements of A instead of just one. In other words, it emphasizes solutions of the system of equations Eq. (2.1)–(2.5). Even though, once one of the components of Ω , say $g(t)$ again, is fixed, the other four will be given by the corresponding universal polynomials of NCSFs in coefficients of $g(t)$ (see Theorem 2.6 below), it is very often not trivial at all what values of these universal polynomials are, or in other words, it is still far away from clear how to write down directly and explicitly the other four components of Ω .

The main aim of this paper is to construct explicitly a NCS system $\Omega_{\mathbb{T}}^W$ over the Grossman-Larson Hopf algebra $\mathcal{H}_{GL}^W$ of W -labeled trees without using any universal polynomials of NCSFs. Once we get the NCS system $\Omega_{\mathbb{T}}^W$ explicitly, then, by Theorem 2.6, these universal polynomials of NCSFs will be transformed into identities of coefficients of the corresponding components of $\Omega_{\mathbb{T}}^W$ (see Remark 4.8). Another immediate consequence is that we also get a very “visualizable representation”, or more formally, a specialization of NCSFs by W -labeled rooted trees, which in return could be useful for studying and understanding certain properties of NCSFs.

Next, let us recall some of the NCSFs first introduced and studied in ([GKLLRT]).

Let $\Lambda = \{\Lambda_m \mid m \geq 1\}$ be a sequence of noncommutative free variables and $\mathcal{NSym}$ or $K\langle\Lambda\rangle$ the free associative algebra generated by Λ over K . For convenience, we also set $\Lambda_0 = 1$. We denote by $\lambda(t)$ the

generating function of Λ_m ($m \geq 0$), i.e. we set

$$(2.6) \quad \lambda(t) := \sum_{m \geq 0} t^m \Lambda_m = 1 + \sum_{k \geq 1} t^k \Lambda_k.$$

In the theory of NCSFs ([GKLLRT]), Λ_m ($m \geq 0$) is the noncommutative analog of the m^{th} classical (commutative) elementary symmetric function and is called the m^{th} (*noncommutative*) *elementary symmetric function*.

To define some other NCSFs, we consider Eqs. (2.2)–(2.5) over the free K -algebra $\mathcal{NSym}$ with $f(t) = \lambda(t)$. The solutions for $g(t)$, $d(t)$, $h(t)$, $m(t)$ exist and are unique, whose coefficients will be the NCSFs that we are going to define. Following the notation in [GKLLRT] and [Z2], we denote the resulted 5-tuple by

$$(2.7) \quad \Pi := (\lambda(t), \sigma(t), \Phi(t), \psi(t), \xi(t))$$

and write the last four generating functions of Π explicitly as follows.

$$(2.8) \quad \sigma(t) = \sum_{m \geq 0} t^m S_m,$$

$$(2.9) \quad \Phi(t) = \sum_{m \geq 1} t^m \frac{\Phi_m}{m}$$

$$(2.10) \quad \psi(t) = \sum_{m \geq 1} t^{m-1} \Psi_m,$$

$$(2.11) \quad \xi(t) = \sum_{m \geq 1} t^{m-1} \Xi_m.$$

Now, for any $m \geq 1$, we define S_m to be the m^{th} (*noncommutative*) *complete homogeneous symmetric function* and Φ_m (resp. Ψ_m) the m^{th} *power sum symmetric function of the second (resp. first) kind*. Note that, Ξ_m ($m \geq 1$) were denoted by Ψ_m^* in [GKLLRT]. Due to Proposition 2.5 below, the NCSFs Ξ_m ($m \geq 1$) do not play an important role in the NCSF theory (see the comments in page 234 in [GKLLRT]). But, in the context of some other problems, relations of Ξ_m 's with other NCSFs, especially, with Ψ_m 's, are also important (see [Z4], for example). So here, following [Z2], we call $\Xi_m \in \mathcal{NSym}$ ($m \geq 1$) the m^{th} (*noncommutative*) *power sum symmetric function of the third kind*.

The following two propositions proved in [GKLLRT] and [KLT] will be very useful for our later arguments.

Proposition 2.4. *For any unital commutative $\mathbb{Q}$ -algebra K , the free algebra $\mathcal{NSym}$ is freely generated by any one of the families of the NCSFs defined above.*

Proposition 2.5. *Let ω_Λ be the anti-involution of $\mathcal{NSym}$ which fixes Λ_m ($m \geq 1$). Then, for any $m \geq 1$, we have*

$$(2.12) \quad \omega_\Lambda(S_m) = S_m,$$

$$(2.13) \quad \omega_\Lambda(\Phi_m) = \Phi_m,$$

$$(2.14) \quad \omega_\Lambda(\Psi_m) = \Xi_m.$$

Next, let us recall the following graded K -Hopf algebra structure of $\mathcal{NSym}$. It has been shown in [GKLLRT] that $\mathcal{NSym}$ is the universal enveloping algebra of the free Lie algebra generated by Ψ_m ($m \geq 1$). Hence, it has a K -Hopf algebra structure as all other universal enveloping algebras of Lie algebras do. Its co-unit $\epsilon : \mathcal{NSym} \rightarrow K$, co-product Δ and antipode S are uniquely determined by

$$(2.15) \quad \epsilon(\Psi_m) = 0,$$

$$(2.16) \quad \Delta(\Psi_m) = 1 \otimes \Psi_m + \Psi_m \otimes 1,$$

$$(2.17) \quad S(\Psi_m) = -\Psi_m,$$

for any $m \geq 1$.

Next, we introduce the *weight* of NCSFs by setting the weight of any monomial $\Lambda_{m_1}^{i_1} \Lambda_{m_2}^{i_2} \cdots \Lambda_{m_k}^{i_k}$ to be $\sum_{j=1}^k i_j m_j$. For any $m \geq 0$, we denote by $\mathcal{NSym}_{[m]}$ the vector subspace of $\mathcal{NSym}$ spanned by the monomials of Λ of weight m . Then it is easy to see that

$$(2.18) \quad \mathcal{NSym} = \bigoplus_{m \geq 0} \mathcal{NSym}_{[m]},$$

which provides a grading for $\mathcal{NSym}$.

Note that, it has been shown in [GKLLRT], for any $m \geq 1$, the NCSFs $S_m, \Phi_m, \Psi_m \in \mathcal{NSym}_{[m]}$. By Proposition 2.5, this is also true for the NCSFs Ξ_m 's. By the facts above and Eqs. (2.15)–(2.17), it is also easy to check that, with the grading given in Eq. (2.18), $\mathcal{NSym}$ forms a graded K -Hopf algebra. Its graded dual is given by the space $\mathcal{QSym}$ of quasi-symmetric functions, which were first introduced by I. Gessel [Ge] (see [MR] and [St2] for more discussions).

Now we come back to our discussions on the NCS systems. From the definitions of the NCSFs above, we see that $(\mathcal{NSym}, \Pi)$ obviously forms a NCS system. More importantly, as shown in Theorem 2.1 in [Z2], we have the following important theorem on the NCS system $(\mathcal{NSym}, \Pi)$.

Theorem 2.6. *Let A be a K -algebra and Ω a NCS system over A . Then,*

(a) *There exists a unique K -algebra homomorphism $\mathcal{S} : \mathcal{NSym} \rightarrow A$ such that $\mathcal{S}^{\times 5}(\Pi) = \Omega$.*

(b) *If A is further a K -bialgebra (resp. K -Hopf algebra) and one of the equivalent statements in Proposition 2.2 holds for the NCS system Ω , then $\mathcal{S} : \mathcal{NSym} \rightarrow A$ is also a homomorphism of K -bialgebras (resp. K -Hopf algebras).*

Remark 2.7. *By applying the similar arguments as in the proof of Theorem 2.6, or simply taking the quotient over the two-sided ideal generated by the commutators of Λ_m 's, it is easy to see that, over the category of commutative K -algebras, the universal NCS system is given by the generating functions of the corresponding classical (commutative) symmetric functions ([Mc]).*

3. The Grossman-Larson Hopf Algebra and The Connes-Kreimer Hopf Algebra

Let K be any unital commutative $\mathbb{Q}$ -algebra and W a non-empty subset of positive integers. In this section, we first fix some notations for unlabeled rooted trees and W -labeled rooted trees that will be used throughout this paper. We then recall the Connes-Kreimer Hopf algebra and the Grossman-Larson Hopf algebra of W -labeled forests and W -labeled rooted trees, respectively. Finally, by using the duality between the Grossman-Larson Hopf algebra and the Connes-Kreimer Hopf algebra (see Theorem 3.2), we prove a technic lemma, Lemma 3.4, that will play an important role in our later arguments.

First, let us fix the following notation which will be used throughout the rest of this paper.

Notation:

By a *rooted tree* we mean a finite 1-connected graph with one vertex designated as its *root*. For convenience, we also view the empty set $\emptyset$ as a rooted tree and call it the *emptyset* rooted tree. The rooted tree with a single vertex is called the *singleton* and denoted by $\circ$. There are natural ancestral relations between vertices. We say a vertex w is a *child* of vertex v if the two are connected by an edge and w lies further from the root than v . In the same situation, we say v is the *parent* of w . A vertex is called a *leaf* if it has no children.

Let $W \subseteq \mathbb{N}^+$ be any non-empty subset of positive integers. A *W -labeled rooted tree* is a rooted tree with each vertex labeled by an element of W . If an element $m \in W$ is assigned to a vertex v , then m

is called the *weight* of the vertex v . When we speak of isomorphisms between unlabeled (resp. W -labeled) rooted trees, we will always mean isomorphisms which also preserve the root (resp. the root and also the labels of vertices). We will denote by $\mathbb{T}$ (resp. $\mathbb{T}^W$) the set of isomorphism classes of all unlabeled (resp. W -labeled) rooted trees. A disjoint union of any finitely many rooted trees (resp. W -labeled rooted trees) is called a *rooted forest* (resp. W -labeled *rooted forest*). We denote by $\mathbb{F}$ (resp. $\mathbb{F}^W$) the set of unlabeled (resp. W -labeled) rooted forests.

With these notions in mind, we establish the following notation.

- (1) For any rooted tree $T \in \mathbb{T}^W$, we set the following notation:
 - rt_T denotes the root vertex of T and $O(T)$ the set of all the children of rt_T . We set $o(T) = |O(T)|$ (the cardinal number of the set $O(T)$).
 - $E(T)$ denotes the set of edges of T .
 - $V(T)$ denotes the set of vertices of T and $v(T) = |V(T)|$.
 - $L(T)$ denotes the set of leaves of T and $l(T) = |L(T)|$.
 - For any $v \in V(T)$, we define the *height* of v to be the number of edges in the (unique) geodesic connecting v to rt_T . The *height* of T is defined to be the maximum of the heights of its vertices.
 - For any $T \in \mathbb{T}^W$ and $T \neq \emptyset$, $|T|$ denotes the sum of the weights of all vertices of T . When $T = \emptyset$, we set $|T| = 0$.
 - For any $T \in \mathbb{T}^W$, we denote by $\text{Aut}(T)$ the automorphism group of T and $\alpha(T)$ the cardinal number of $\text{Aut}(T)$.
- (2) Any subset of $E(T)$ is called a *cut* of T . A cut $C \subseteq E(T)$ is said to be *admissible* if no two different edges of C lie in the path connecting the root and a leaf. We denote by $\mathcal{C}(T)$ the set of all admissible cuts of T . Note that, the empty subset $\emptyset$ of $E(T)$ and $C = \{e\}$ for any $e \in E(T)$ are always admissible cuts. We will identify any edge $e \in E(T)$ with the admissible cut $C := \{e\}$ and simply say the edge e itself is an admissible cut of T .
- (3) For any $T \in \mathbb{T}^W$ with $T \neq \circ$, let $C \in \mathcal{C}(T)$ be an admissible cut of T with $|C| = m \geq 1$. Note that, after deleting the edges in C from T , we get a disjoint union of $m+1$ rooted trees, say $T_0, T_1, \dots, T_m$ with $\text{rt}(T) \in V(T_0)$. We define $R_C(T) = T_0 \in \mathbb{T}^W$ and $P_C(T) \in \mathbb{F}^W$ the rooted forest formed by $T_1, \dots, T_m$.
- (4) For any disjoint admissible cuts C_1 and C_2 , we say “ C_1 lies above C_2 ”, and write $C_1 \succ C_2$, if $C_2 \subseteq E(R_{C_1}(T))$. This merely says that all edges of C_2 remain when we remove all edges of C_1 and $P_{C_1}(T)$. Note that this relation is not transitive. When we

write $C_1 \succ \cdots \succ C_r$ for $C_1, \dots, C_r \in \mathcal{C}(T)$, we will mean that $C_i \succ C_j$ whenever $i < j$.

- (5) For any $T \in \mathbb{T}^W$, we say T is a *chain* if its underlying rooted tree is a rooted tree with a single leaf. We say T is a *shrub* if its underlying rooted tree is a rooted tree of height 1. We say T is *primitive* if its root has only one child. For any $m \geq 1$, we set $\mathbb{H}_m$, $\mathbb{S}_m$ and $\mathbb{P}_m$ to be the sets of the chains, shrubs and primitive rooted trees T of weight $|T| = m$, respectively. $\mathbb{H}$, $\mathbb{S}$ and $\mathbb{P}$ are set to be the unions of $\mathbb{H}_m$, $\mathbb{S}_m$ and $\mathbb{P}_m$, respectively, for all $m \geq 1$.

For example, in the case where $W = \{1\}$, which allows not to write the labels, we have

$$\begin{aligned} \mathbb{H} &= \left\{ \cdot, \text{1}, \text{1}, \text{1}, \text{1}, \text{1} \dots \right\}, \\ \mathbb{S} &= \left\{ \text{1}, \text{V}, \text{V}, \text{V}, \text{V} \dots \right\}, \\ \mathbb{P} &= \left\{ \text{1}, \text{1}, \text{Y}, \text{1}, \text{Y}, \text{Y}, \text{Y}, \text{1}, \text{1} \dots \right\}. \end{aligned}$$

For any non-empty $W \subseteq \mathbb{N}^+$, we define the following operations for W -labeled rooted forests. For any $F \in \mathbb{F}^W$ which is disjoint union of W -labeled rooted trees T_i ($1 \leq i \leq m$), we set $B_+(T_1, T_2, \dots, T_m)$ the rooted tree obtained by connecting roots of T_i ($1 \leq i \leq m$) to a newly added root. We will keep the labels for the vertices of $B_+(T_1, T_2, \dots, T_m)$ from T_i 's, but for the root, we label it by 0. For convenience, we also fix the following short convention for the operation B_+ . For the empty rooted tree $\emptyset$, we set $B_+(\emptyset)$ to be the singleton labeled by 0. For any $T_i \in \mathbb{T}^W$ ($1 \leq i \leq m$) and $j_i \geq 1$, the notation $B_+(T_1^{j_1}, T_2^{j_2}, \dots, T_m^{j_m})$ denotes the rooted tree obtained by applying the operation B_+ to j_1 -copies of T_1 ; j_2 -copies of T_2 ; and so on. Later, for any unital $\mathbb{Q}$ -algebra K and $m \geq 1$, we will also extend the operation B_+ multi-linearly to a linear map B_+ from $(\mathcal{H}_{CK}^W)^{\times m}$ to $\mathcal{H}_{GL}^W$, where $\mathcal{H}_{CK}^W$ and $\mathcal{H}_{GL}^W$ at this moment are the vector spaces spanned over K by the elements of $\mathbb{T}^W$ and $B_+(\mathbb{T}^W)$, respectively.

Now, we set $\bar{\mathbb{T}}^W := \{B_+(F) \mid F \in \mathbb{F}^W\}$. Then, $B_+ : \mathbb{F}^W \rightarrow \bar{\mathbb{T}}^W$ becomes a bijection. We denote by $B_- : \bar{\mathbb{T}}^W \rightarrow \mathbb{F}^W$ the inverse map of B_+ . More precisely, for any $T \in \bar{\mathbb{T}}^W$, $B_-(T)$ is the W -labeled rooted forest obtained by cutting off the root of T as well as all edges connecting to the root in T .

Note that, precisely speaking, elements of $\bar{\mathbb{T}}^W$ are not W -labeled trees for $0 \notin W$. But, if we set $\bar{W} = W \cup \{0\}$, then we can view $\bar{\mathbb{T}}^W$ as a subset of $\bar{W}$ -labeled rooted trees T with the root rt_T labeled by 0 and all other vertices labeled by non-zero elements of $\bar{W}$. We extend the definition of the weight for elements of $\mathbb{F}^W$ to elements of $\bar{\mathbb{T}}^W$ by simply counting the weight of roots by zero. We set $\bar{\mathbb{S}}_m^W := B_+(\mathbb{S}_m^W)$ ($m \geq 1$) and $\bar{\mathbb{S}}^W := B_+(\mathbb{S}^W)$. We also define $\bar{\mathbb{H}}_m^W, \bar{\mathbb{P}}_m^W, \bar{\mathbb{H}}^W$ and $\bar{\mathbb{P}}^W$ in the similar way.

Next we fix a unital commutative $\mathbb{Q}$ -algebra K and a non-empty subset of positive integers W , and first recall the Connes-Kreimer Hopf algebras $\mathcal{H}_{CK}^W$ of W -labeled rooted forests.

As a K -algebra, the Connes-Kreimer Hopf algebra $\mathcal{H}_{CK}^W$ is the free commutative algebra generated by formal variables $\{X_T | T \in \mathbb{T}^W\}$. Here, for convenience, we will still use T to denote the variable X_T in $\mathcal{H}_{CK}^W$. The K -algebra product is given by the disjoint union. The identity element of this algebra, denoted by 1, is the free variable $X_\emptyset$ corresponding to the emptyset rooted tree $\emptyset$. The coproduct $\Delta : \mathcal{H}_{CK}^W \rightarrow \mathcal{H}_{CK}^W \otimes \mathcal{H}_{CK}^W$ is uniquely determined by setting

$$(3.1) \quad \Delta(1) = 1 \otimes 1,$$

$$(3.2) \quad \Delta(T) = T \otimes 1 + \sum_{C \in \mathcal{C}(T)} P_C(T) \otimes R_C(T).$$

The co-unit $\epsilon : \mathcal{H}_{CK}^W \rightarrow K$ is the K -algebra homomorphism which sends $1 \in \mathcal{H}_{CK}^W$ to $1 \in K$ and T to 0 for any $T \in \mathbb{T}^W$ with $T \neq \emptyset$. With the operations defined above and the grading given by the weight, the vector space $\mathcal{H}_{CK}^W$ forms a connected graded commutative bialgebra. Since any connected graded bialgebra is a Hopf algebra, there is a unique antipode $S : \mathcal{H}_{CK}^W \rightarrow \mathcal{H}_{CK}^W$ that makes $\mathcal{H}_{CK}^W$ a connected graded commutative K -Hopf algebra. For a formula for the antipode, see [F].

Next we recall the Grossman-Larson Hopf algebra of labeled rooted trees. As a vector space, the Grossman-Larson Hopf algebra $\mathcal{H}_{GL}^W$ is the vector space spanned by elements of $\bar{\mathbb{T}}^W$ over K . For any $T \in \bar{\mathbb{T}}^W$, we will still denote by T the vector in $\mathcal{H}_{GL}^W$ that is corresponding to T . The algebra product is defined as follows.

For any $T, S \in \bar{\mathbb{T}}^W$ with $T = B_+(T_1, T_2, \dots, T_m)$, we set $T \cdot S$ to be the sum of the rooted trees obtained by connecting the roots of T_i ($1 \leq i \leq m$) to vertices of S in all possible $m^{v(S)}$ different ways. Note that, the identity element with respect to this algebra product is given by the singleton $\circ = B_+(\emptyset)$. But we will denote it by 1.

To define the co-product $\Delta : \mathcal{H}_{GL}^W \rightarrow \mathcal{H}_{GL}^W \otimes \mathcal{H}_{GL}^W$, we first set

$$(3.3) \quad \Delta(\circ) = \circ \otimes \circ.$$

Now let $T \in \bar{\mathbb{T}}^W$ with $T \neq \circ$, say $T = B_+(T_1, T_2, \dots, T_m)$ with $m \geq 1$ and $T_i \in \mathbb{T}^W$ ($1 \leq i \leq m$). For any non-empty subset $I \subseteq \{1, 2, \dots, m\}$, we denote by $B_+(T_I)$ the rooted tree obtained by applying the B_+ operation to the rooted trees T_i with $i \in I$. For convenience, when $I = \emptyset$, we set $B_+(T_I) = 1$. With this notation fixed, the co-product for T is given by

$$(3.4) \quad \Delta(T) = \sum_{I \sqcup J = \{1, 2, \dots, m\}} B_+(T_I) \otimes B_+(T_J).$$

The co-unit $\epsilon : \mathcal{H}_{GL}^W \rightarrow K$ is the K -algebra homomorphism which sends $1 \in \mathcal{H}_{GL}^W$ to $1 \in K$ and T to 0 for any $T \in \bar{\mathbb{T}}^W$ with $T \neq \emptyset$. With the operations defined above and the grading given by the weight, the vector space $\mathcal{H}_{GL}^W$ forms a graded commutative bialgebra. Therefore, there is a unique antipode $S : \mathcal{H}_{CK}^W \rightarrow \mathcal{H}_{CK}^W$ that makes $\mathcal{H}_{CK}^W$ a graded K -Hopf algebra. Actually, by the general recurrent formula, we can write down the antipode of $\mathcal{H}_{GL}^W$ as follows.

Note that the singleton $\circ$ is a group-like element and $S(\circ) = \circ^{-1} = \circ$. Now assume $T \neq \circ$ and write $T = B_+(T_1, T_2, \dots, T_m)$ with $m \geq 1$ and $T_i \in \mathbb{T}^W$. Let $I := \{1, 2, \dots, m\}$. For $1 \leq r \leq m$, let $\mathcal{P}_r$ be the set of r -tuples $(I_1, I_2, \dots, I_r)$ of disjoint non-empty subsets of I whose union is I . In other words, $\mathcal{P}_r$ is the set of all ordered partitions of I into r non-empty subsets of I .

Lemma 3.1. *Let S denote the antipode of the Grossman-Larson Hopf algebra $\mathcal{H}_{GL}^W$ of W -labeled rooted trees. Then, for any $T \in \bar{\mathbb{T}}^W$ with $T \neq \circ$ as above, we have*

$$(3.5) \quad S(T) = \sum_{r=1}^m (-1)^r \sum_{(I_1, \dots, I_r) \in \mathcal{P}_r} B_+(T_{I_1}) B_+(T_{I_2}) \cdots B_+(T_{I_r})$$

Proof: By the general recurrent formula for the antipode of connected cocommutative graded Hopf algebras, we know that the antipode S of $\mathcal{H}_{GL}^W$ satisfies the following equation:

$$(3.6) \quad S(T) = -T - \sum_{(I, J) \in \mathcal{P}_2} S(B_+(T_I)) B_+(T_J).$$

Then it is easy to check directly that, for any $T \in \bar{\mathbb{T}}^W$, $S(T)$ given by Eq. (3.5) does satisfy Eq. (3.6). Since the solution to Eq. (3.6) is unique, the antipode S of $\mathcal{H}_{GL}^W$ is actually given by Eq. (3.5). $\square$

Note that, from Eq. (3.4), it is easy to see that, a rooted tree $T \in \bar{\mathbb{T}}^W$ is a primitive element of the Hopf algebra $\mathcal{H}_{GL}^W$ iff it is a primitive rooted tree in the sense that we defined before, namely the root of T has one

and only one child. It is noticeable that the set of primitive W -labeled rooted trees is a basis of the space $\text{Prim}(\mathcal{H}_{GL}^W)$ of primitive elements of $\mathcal{H}_{GL}^W$. Moreover, by the Milnor-Moore theorem, $\mathcal{H}_{GL}^W$ is isomorphic to $\mathcal{U}(\text{Prim}(\mathcal{H}_{GL}^W))$.

The relation between the Grossman-Larson Hopf algebra $\mathcal{H}_{GL}^W$ and the Connes-Kreimer Hopf algebra $\mathcal{H}_{CK}^W$ is given by the following important theorem, which was proved in [H] and [F].

Theorem 3.2. *The Hopf algebras $\mathcal{H}_{GL}^W$ and $\mathcal{H}_{CK}^W$ are graded dual to each other. The pairing is given by, for any $T \in \bar{\mathbb{T}}^W$ and $S \in \mathbb{F}^W$,*

$$(3.7) \quad \langle T, F \rangle = \begin{cases} 0, & \text{if } T \not\simeq B_+(F), \\ \alpha(T), & \text{if } T \simeq B_+(F). \end{cases}$$

Furthermore, the following theorem on the algebra structure constants of $\mathcal{H}_{GL}^W$ was also proved in [H] and [F].

Theorem 3.3. *For any $T', S \in \bar{\mathbb{T}}^W$, We have*

$$(3.8) \quad T' \cdot S = \sum_{T \in \bar{\mathbb{T}}^W} \sum_{\substack{C \in \mathcal{C}(T) \\ B_+(P_C(T)) \sim T', \\ R_C(T) \sim S}} \frac{\alpha(T')\alpha(S)}{\alpha(T)} T.$$

Note that, Eq. (3.8) suggests that it is much more convenient to work with the basis $\{\mathcal{V}_T := T/\alpha(T) \mid T \in \bar{\mathbb{T}}^W\}$ than the basis $\{T \mid T \in \bar{\mathbb{T}}^W\}$. For example, in terms of $\mathcal{V}_T$, Eq. (3.8) becomes

$$(3.9) \quad \mathcal{V}_{T'} \cdot \mathcal{V}_S = \sum_{T \in \bar{\mathbb{T}}^W} \sum_{\substack{C \in \mathcal{C}(T) \\ B_+(P_C(T)) \sim T', \\ R_C(T) \sim S}} \mathcal{V}_T.$$

Finally, we extend Theorem 3.3 to a more general setting (see Lemma 3.4 below). It can be viewed as a generalization of Lemma 2.8 in [WZ] which essentially is the case of Lemma 3.4 when only primitive rooted trees are involved. First, let us fix the following notation.

Let $\vec{C} = (C_1, \dots, C_r) \in \mathcal{C}(T)^{\times r}$ be a sequence of admissible cuts with $C_1 \succ \dots \succ C_r$. We define a sequence of $T_{\vec{C},1}, \dots, T_{\vec{C},r+1} \in \bar{\mathbb{T}}^W$ as follows: we first set $T_{\vec{C},1} = B_+(P_{C_1}(T))$ and let $S_1 = R_{C_1}(T)$. Note that $C_2, \dots, C_r \in \mathcal{C}(S_1)$. We then set $T_{\vec{C},2} = B_+(P_{C_2}(S_1))$ and $S_2 = R_{C_2}(S_1)$ and repeat this procedure until we get $S_r = R_{C_r}(S_{r-1})$ and then set $T_{\vec{C},r+1} = S_r$. In the case that, each C_i ($1 \leq i \leq r$) consists of a single edge, say $e_i \in E(T)$, we simply denote $T_{\vec{C},i}$ by T_{e_i} .

Now we fix a positive integer r and let $y = \{y_T^{(i)} \mid 1 \leq i \leq r; T \in \bar{\mathbb{T}}^W\}$ be a collection of commutative formal variables.

Lemma 3.4. *For any $r \geq 1$, we have,*

$$(3.10) \quad \sum_{(T_1, \dots, T_r) \in (\bar{\mathbb{T}}^W)^r} \left[y_{T_1}^{(1)} \mathcal{V}_{T_1} \right] \cdots \left[y_{T_r}^{(r)} \mathcal{V}_{T_r} \right] \\ = \sum_{T \in \bar{\mathbb{T}}^W} \sum_{\substack{\vec{C} = (C_1, \dots, C_{r-1}) \in \mathcal{C}(T)^{r-1} \\ C_1 \succ \dots \succ C_{r-1}}} y_{T_{\vec{C},1}}^{(1)} \cdots y_{T_{\vec{C},r}}^{(r)} \mathcal{V}_T.$$

Proof: We denote the LHS of Eq. (3.10) by Q and write it as

$$Q = \sum_{T \in \bar{\mathbb{T}}^W} y_T \mathcal{V}_T.$$

Then, by Eq. (3.7) $y_T = \langle Q, B_-(T) \rangle$ for any $T \in \bar{\mathbb{T}}^W$, where $B_-(T)$ is the forest obtained by deleting the root of T . So:

$$y_T = \sum_{(T_1, \dots, T_r)} y_{T_1}^{(1)} \cdots y_{T_r}^{(r)} \langle \mathcal{V}_{T_1} \cdots \mathcal{V}_{T_r}, B_-(T) \rangle \\ = \sum_{(T_1, \dots, T_r)} y_{T_1}^{(1)} \cdots y_{T_r}^{(r)} \langle \mathcal{V}_{T_1} \otimes (\mathcal{V}_{T_2} \cdots \mathcal{V}_{T_r}), \Delta(B_-(T)) \rangle \\ = \sum_{(T_1, \dots, T_r)} y_{T_1}^{(1)} \cdots y_{T_r}^{(r)} \langle \mathcal{V}_{T_1} \otimes \mathcal{V}_{T_2} \otimes (\mathcal{V}_{T_3} \cdots \mathcal{V}_{T_r}), (I \otimes \Delta) \circ \Delta(B_-(T)) \rangle,$$

where I is the identity map of $\mathcal{H}_{CK}^W$. Repeating the process above:

$$= \sum_{(T_1, \dots, T_r)} y_{T_1}^{(1)} \cdots y_{T_r}^{(r)} \langle \mathcal{V}_{T_1} \otimes \cdots \otimes \mathcal{V}_{T_r}, (I^{\otimes(r-2)} \otimes \Delta) \circ \cdots \circ \Delta(B_-(T)) \rangle.$$

One the other hand, by definition of the coproduct of $\mathcal{H}_{CK}^W$ and definition of $\succ$, we have

$$(I^{\otimes(r-2)} \otimes \Delta) \circ \cdots \circ (I \otimes \Delta) \circ \Delta(B_-(T)) \\ = \sum_{\substack{\vec{C} = (C_1, \dots, C_{r-1}) \in \mathcal{C}(T)^{r-1} \\ C_1 \succ \dots \succ C_{r-1}}} B_-(T_{\vec{C},1}) \otimes \cdots \otimes B_-(T_{\vec{C},r}).$$

Therefore, we get

$$y_T = \sum_{\substack{\vec{C} = (C_1, \dots, C_{r-1}) \in \mathcal{C}(T)^{r-1} \\ C_1 \succ \dots \succ C_{r-1}}} y_{T_{\vec{C},1}}^{(1)} \cdots y_{T_{\vec{C},r}}^{(r)}.$$

□

4. A NCS System over the Grossman-Larson Hopf Algebra $\mathcal{H}_{GL}^W$ of W -Labeled Rooted Trees

In this section, for any non-empty $W \subseteq \mathbb{N}^+$, we construct a NCS system $\Omega_{\mathbb{T}}^W$ over the Grossman-Larson Hopf algebra $\mathcal{H}_{GL}^W$. First, let us introduce the following generating functions of certain elements of $\mathcal{H}_{GL}^W$, which will be the components of the NCS system $\Omega_{\mathbb{T}}^W$ corresponding to $f(t)$, $g(t)$, $h(t)$ and $m(t)$ according the notation in Definition 2.1.

$$(4.1) \quad \tilde{f}(t) := \sum_{T \in \mathbb{S}^W} (-1)^{o(T)+|T|} t^{|T|} \mathcal{V}_T = 1 + \sum_{\substack{T \in \mathbb{S}^W \\ T \neq \circ}} (-1)^{o(T)+|T|} t^{|T|} \mathcal{V}_T,$$

$$(4.2) \quad \tilde{g}(t) := \sum_{T \in \mathbb{T}^W} t^{|T|} \mathcal{V}_T = 1 + \sum_{\substack{T \in \mathbb{T}^W \\ T \neq \circ}} t^{|T|} \mathcal{V}_T,$$

$$(4.3) \quad \tilde{h}(t) := \sum_{T \in \mathbb{H}^W} t^{|T|-1} \beta_T \mathcal{V}_T,$$

$$(4.4) \quad \tilde{m}(t) := \sum_{T \in \mathbb{P}^W} t^{|T|-1} \gamma_T \mathcal{V}_T,$$

where, for any $T \in \mathbb{H}^W$ (resp. $T \in \mathbb{P}^W$), β_T (resp. γ_T) is the weight of the unique leaf (resp. the unique child of the root) of T . Note that, for the singleton $T = \circ$, we have $\beta_T = \gamma_T = 0$. So $\tilde{h}(0) = \tilde{m}(0) = 0$.

For example, when $W = \{1, 2\}$, we have

$$\begin{aligned} \tilde{f}(t) &= 1 + \mathcal{V}_{\mathbf{1}^1} t + \left(\mathcal{V}_{\mathbf{1}_{\mathbf{V}^1}} - \mathcal{V}_{\mathbf{1}^2} \right) t^2 + \left(\mathcal{V}_{\mathbf{1}_{\mathbf{V}^1}^1} - \mathcal{V}_{\mathbf{1}_{\mathbf{V}^2}} \right) t^3 + \dots \\ \tilde{g}(t) &= 1 + \mathcal{V}_{\mathbf{1}^1} t + \left(\mathcal{V}_{\mathbf{1}^2} + \mathcal{V}_{\mathbf{1}_{\mathbf{V}^1}} + \mathcal{V}_{\mathbf{1}_{\mathbf{1}}^1} \right) t^2 \\ &\quad + \left(\mathcal{V}_{\mathbf{1}_{\mathbf{V}^2}} + \mathcal{V}_{\mathbf{1}_{\mathbf{1}}^2} + \mathcal{V}_{\mathbf{1}_{\mathbf{2}}^1} + \mathcal{V}_{\mathbf{1}_{\mathbf{V}^1}^1} + \mathcal{V}_{\mathbf{1}_{\mathbf{V}^1}^1} + \mathcal{V}_{\mathbf{1}_{\mathbf{V}^1}^1} + \mathcal{V}_{\mathbf{1}_{\mathbf{1}}^1} \right) t^3 + \dots \\ \tilde{h}(t) &= \mathcal{V}_{\mathbf{1}^1} + \left(\mathcal{V}_{\mathbf{1}_{\mathbf{1}}^1} + 2\mathcal{V}_{\mathbf{1}^2} \right) t + \left(\mathcal{V}_{\mathbf{1}_{\mathbf{1}}^1} + 2\mathcal{V}_{\mathbf{1}_{\mathbf{1}}^2} + \mathcal{V}_{\mathbf{1}_{\mathbf{2}}^1} \right) t^2 + \dots \\ \tilde{m}(t) &= \mathcal{V}_{\mathbf{1}^1} + \left(\mathcal{V}_{\mathbf{1}_{\mathbf{1}}^1} + 2\mathcal{V}_{\mathbf{1}^2} \right) t + \left(\mathcal{V}_{\mathbf{1}_{\mathbf{1}}^2} + 2\mathcal{V}_{\mathbf{1}_{\mathbf{2}}^1} + \mathcal{V}_{\mathbf{1}_{\mathbf{V}^1}^1} + \mathcal{V}_{\mathbf{1}_{\mathbf{1}}^1} \right) t^2 + \dots \end{aligned}$$

Note that, from Eq.(4.1), we have

$$(4.5) \quad \tilde{f}(-t) = \sum_{T \in \mathbb{S}^W} (-1)^{o(T)} t^{|T|} \mathcal{V}_T = 1 + \sum_{\substack{T \in \mathbb{S}^W \\ T \neq \circ}} (-1)^{o(T)} t^{|T|} \mathcal{V}_T.$$

A different way to look at the generating function $\tilde{f}(-t)$ is as follows. For any $m \in W$, let κ_m denote the singleton labeled by m and set

$$(4.6) \quad \kappa(t) := \sum_{m \in W} t^m \kappa_m.$$

Lemma 4.1.

$$(4.7) \quad \tilde{f}(-t) = 1 + \sum_{d \geq 1} \frac{(-1)^d}{d!} B_+(\kappa(t)^d),$$

where $B_+(\kappa(t)^d)$ denotes the element obtained by applying B_+ to d -copies of $\kappa(t)$.

Proof: First, it is easy to see that, the only terms that can appear in the expansion of the RHS of Eq. (4.7) are shrubs. Secondly, from the definition of the operation B_+ , we see that B_+ is symmetric and multilinear in its components. Therefore, we can expand the term $B_+(\kappa(t)^d)$ into a linear combination of rooted trees $S \in \mathbb{T}^W$ in a similar way as we expand the power $(\sum_{m \in W} t^m u_m)^d$ for some free commutative variables u_m ($m \in W$).

Now, for any shrub $S \in \bar{\mathbb{S}}^W$ with $S \neq \circ$, let $\{m_j \in W \mid 1 \leq j \leq N\}$ be the set of all labels of the leaves of S . Let $i_j \geq 1$ ($1 \leq j \leq N$) be the number of the m_j -labeled leaves of S . Then we have

$$(4.8) \quad o(S) = \sum_{1 \leq j \leq N} i_j,$$

$$(4.9) \quad |S| = \sum_{1 \leq j \leq N} i_j m_j,$$

$$(4.10) \quad \alpha(S) = \prod_{1 \leq j \leq N} (i_j)!.$$

Now let us consider the coefficient c_S of S in the linear expansion of the RHS of Eq. (4.7). By the observations in the first paragraph of the proof and Eqs. (4.8)-(4.10), it is easy to see that we have

$$\begin{aligned} c_S &= \frac{(-1)^{o(S)} t^{|S|}}{o(S)!} \binom{o(S)}{i_1, \dots, i_N} \\ &= (-1)^{o(S)} \frac{t^{|S|}}{\prod_{1 \leq j \leq N} (i_j)!} \\ &= (-1)^{o(S)} \frac{t^{|S|}}{\alpha(S)}, \end{aligned}$$

which is same as the coefficient of S in $\tilde{f}(-t)$ since $\mathcal{V}_S = \frac{1}{\alpha(S)} S$. Hence we are done. $\square$

To define the generating function $\tilde{d}(t)$ for the third component of the under-construction NCS system $\Omega_{\mathbb{T}}^W$, we first need the following definition.

Definition 4.2. (a) We define a constant $\theta_T \in \mathbb{Q}$ for each unlabeled rooted tree T as follows.

- (1) For the singleton $\circ$ and any non-primitive rooted tree $T \in \mathbb{T}$, i.e. $o(T) > 1$, we set $\theta_{\circ} = \theta_T = 0$.
- (2) For $T = B_+(\circ)$, we set $\theta_T = 1$.
- (3) For any primitive $T \in \mathbb{P}$ with $v(T) \geq 3$, we define θ_T inductively by

$$(4.11) \quad \theta_T = 1 - \sum_{m \geq 2} \frac{1}{m!} \sum_{\substack{\vec{e} = (e_1, \dots, e_{m-1}) \in E(T)^{m-1} \\ e_1 \succ \dots \succ e_{m-1}}} \theta_{T_{e_1}} \theta_{T_{e_2}} \cdots \theta_{T_{e_m}},$$

where T_{e_i} 's in the equation above have been defined before Lemma 3.4.

(b) For any $W \subseteq \mathbb{N}$ and W -labeled rooted tree T , we set $\theta_T := \theta_{\bar{T}}$, where $\bar{T}$ is the underlying unlabeled rooted tree of T .

Remark 4.3. As we will show later in Section 5, the constant θ_T ($T \in \mathbb{T}$) has a natural combinatorial interpretations as follows. If we write $T = B_+(F)$ for some rooted forest $F \in \mathbb{F}$ and let $\Omega(F, s)$ be the order polynomial (see [St1]) of F , then θ_T will be the coefficient of s of $\Omega(F, s)$ (see Proposition 5.8 in Section 5). Furthermore, if we denote by $\nabla : K[s] \rightarrow K[s]$ the linear operator which maps any $f(s) \in K[s]$ to $f(s) - f(s-1)$, then θ_T is also the coefficient of s of the polynomial $\nabla \Omega(T, s)$ (see Corollary 5.9). In order to keep our on-going arguments more focus, we will postpone to Section 5 a detailed discussion on these combinatorial interpretations of θ_T ($T \in \mathbb{T}$).

Example 4.4. By Eqs. (5.3), (5.5), (5.10) and (5.13) in Section 5, it is easy to check that, for the chains C_m 's and $B_+(S_m)$ of the shrubs S_m 's, we have

$$(4.12) \quad \theta_{C_m} = \frac{1}{m-1} \quad \text{for any } m \geq 2.$$

$$(4.13) \quad \theta_{B_+(S_m)} = (-1)^m b_m \quad \text{for any } m \geq 0,$$

where b_m ($m \geq 0$) are the Bernoulli numbers which are defined by the generating function

$$(4.14) \quad \frac{x}{e^x - 1} = \sum_{m=0}^{\infty} b_m \frac{x^m}{m!}.$$

Now, we introduce the following generating function:

$$(4.15) \quad \tilde{d}(t) := \sum_{T \in \mathbb{P}^W} t^{|T|} \theta_T \mathcal{V}_T.$$

For example, when $W = \{1, 2\}$, we have

$$\begin{aligned} \tilde{d}(t) = & \mathcal{V}_{\mathbf{1}^1} t + \left(\frac{1}{2} \mathcal{V}_{\mathbf{1}^1_1} + \mathcal{V}_{\mathbf{1}^2} \right) t^2 \\ & + \left(\frac{1}{2} \mathcal{V}_{\mathbf{1}^2_1} + \frac{1}{2} \mathcal{V}_{\mathbf{1}^1_2} + \frac{1}{6} \mathcal{V}_{\mathbf{1}^1_1} + \frac{1}{3} \mathcal{V}_{\mathbf{1}^1_1} \right) t^3 + \dots \end{aligned}$$

Set

$$(4.16) \quad \Omega_{\mathbb{T}}^W := (\tilde{f}(t), \tilde{g}(t), \tilde{d}(t), \tilde{h}(t), \tilde{m}(t)).$$

Then, the main result of this section is the following theorem.

Theorem 4.5. *For any non-empty set $W \subseteq \mathbb{N}$, $\Omega_{\mathbb{T}}^W$ forms a NCS system over the Grossman-Larson Hopf algebra $\mathcal{H}_{GL}^W$.*

Proof: Note that, by Eqs. (4.1) and (4.2), we have $\tilde{f}(0) = \tilde{g}(0) = 1$, hence it will be enough to show Eqs. (2.2)–(2.5) in Definition 2.1 are satisfied by the generating functions in $\Omega_{\mathbb{T}}^W$.

Let us start with Eq. (2.2). First, note that, since $\tilde{g}(0) = 1$, $\tilde{g}(t)$ as an element of $\mathcal{H}_{GL}^W[[t]]$ does have both left and right inverses. So we only need show $\tilde{f}(-t)\tilde{g}(t) = 1$ for $\tilde{g}(t)\tilde{f}(-t) = 1$ will follow automatically. Secondly, by Eq. (4.5) and Lemma 3.4 with $y_T^{(1)} = (-1)^{o(T)}t^{|T|}$ if $T \in \bar{\mathbb{S}}^W$ and 0 otherwise, and $y_T^{(2)} = t^{|T|}$ for any $T \in \bar{\mathbb{T}}^W$, we have

$$\begin{aligned} (4.17) \quad \tilde{f}(-t)\tilde{g}(t) &= \left(\sum_{T' \in \bar{\mathbb{S}}^W} (-1)^{o(T')} t^{|T'|} \mathcal{V}_{T'} \right) \left(\sum_{T'' \in \bar{\mathbb{T}}^W} t^{|T''|} \mathcal{V}_{T''} \right) \\ &= \sum_{T \in \bar{\mathbb{T}}^W} t^{|T|} \left(\sum_{\substack{C \in \mathcal{C}(T) \\ B_+(P_C(T)) \in \bar{\mathbb{S}}^W}} (-1)^{o(B_+(P_C(T)))} \right) \mathcal{V}_T \end{aligned}$$

First, note that, for any rooted tree T and an admissible cut C of T , $B_+(P_C(T))$ is a shrub iff each edge in C is the unique edge connecting with a leaf of T . Therefore, the set of all admissible cuts C such that $B_+(P_C(T)) \in \bar{\mathbb{S}}^W$ is in 1-1 correspondence with the set of subsets of leaves of T . Secondly, when $B_+(P_C(T)) \in \bar{\mathbb{S}}^W$ for an admissible cut C , $o(B_+(P_C(T)))$ is same as the cardinal number $|C|$ of the cut C . With

these observations, for any $T \in \bar{\mathbb{T}}^W$ with $l(T) := |L(T)| > 0$, we have

$$(4.18) \quad \sum_{\substack{C \in \mathcal{C}(T) \\ B_+(P_C(T)) \in \bar{\mathbb{S}}^W}} (-1)^{|C|} = \sum_{k=0}^{l(T)} (-1)^k \binom{l(T)}{k} = 0.$$

Combining Eqs. (4.17) and (4.18), we get $\tilde{f}(-t)\tilde{g}(t) = 1$ and hence Eq. (2.2) for the system $\Omega_{\mathbb{T}}^W$.

Now, let us prove Eq. (2.3) as follows.

$$\begin{aligned} e^{\tilde{d}(t)} &= \sum_{k \geq 0} \frac{1}{k!} \tilde{d}(t)^k \\ &= 1 + \sum_{k \geq 1} \frac{1}{k!} \left(\sum_{T \in \bar{\mathbb{P}}^W} t^{|T|} \theta_T \mathcal{V}_T \right)^k \end{aligned}$$

Applying Lemma 3.4:

$$\begin{aligned} &= 1 + \sum_{T \in \bar{\mathbb{P}}^W} t^{|T|} \theta_T \mathcal{V}_T \\ &\quad + \sum_{k \geq 2} \frac{1}{k!} \sum_{T \in \bar{\mathbb{T}}^W} t^{|T|} \left(\sum_{\substack{\vec{e} = (e_1, \dots, e_{k-1}) \in E(T)^{k-1} \\ e_1 \succ \dots \succ e_{k-1}}} \theta_{T_{\vec{e},1}} \cdots \theta_{T_{\vec{e},k}} \right) \mathcal{V}_T \\ &= 1 + \sum_{T \in \bar{\mathbb{T}}^W} t^{|T|} \left(\theta_T + \sum_{k \geq 2} \frac{1}{k!} \sum_{\substack{\vec{e} = (e_1, \dots, e_{k-1}) \in E(T)^{k-1} \\ e_1 \succ \dots \succ e_{k-1}}} \theta_{T_{\vec{e},1}} \cdots \theta_{T_{\vec{e},k}} \right) \mathcal{V}_T \end{aligned}$$

Applying Eq. (4.11):

$$\begin{aligned} &= 1 + \sum_{T \in \bar{\mathbb{T}}^W} t^{|T|} \mathcal{V}_T \\ &= \tilde{g}(t). \end{aligned}$$

Therefore, we get $e^{\tilde{d}(t)} = \tilde{g}(t)$, which is Eq. (2.3) for the system $\Omega_{\mathbb{T}}^W$.

To prove Eq. (2.4), first, by Lemma 3.4, we have

$$(4.19) \quad \tilde{m}(t)\tilde{g}(t) = \left(\sum_{T' \in \bar{\mathbb{P}}^W} \gamma(T') t^{|T'|-1} \mathcal{V}_{T'} \right) \left(\sum_{T'' \in \bar{\mathbb{T}}^W} t^{|T''|} \mathcal{V}_{T''} \right)$$

$$\begin{aligned}
&= \sum_{T \in \bar{\mathbb{T}}^W} t^{|T|-1} \left(\sum_{\substack{C \in \mathcal{C}(T) \\ B_+(P_C(T)) \in \bar{\mathbb{P}}^W}} \gamma(B_+(P_C(T))) \right) \mathcal{V}_T \\
&= \sum_{T \in \bar{\mathbb{T}}^W} t^{|T|-1} \left(\sum_{e \in E(T)} \gamma(B_+(P_e(T))) \right) \mathcal{V}_T,
\end{aligned}$$

where the last equality follows from the fact that, for any $C \in \mathcal{C}(T)$, $o(B_+(P_C(T))) = 1$ iff C consists of a single edge.

Note that, for any $e \in E(T)$, $\gamma(B_+(P_e(T))) = wt(v'_e)$, where v'_e is the vertex of e which is further away from the root of T . Therefore, continuing with the equation above, we have

$$\tilde{m}(t)\tilde{g}(t) = \sum_{T \in \bar{\mathbb{T}}^W} |T| t^{|T|-1} \mathcal{V}_T = \frac{d\tilde{g}(t)}{dt}.$$

Hence, we get Eq. (2.4) for the system $\Omega_{\mathbb{T}}^W$.

Finally, we show Eq. (2.5). First, by Lemma 3.4, we have

$$\begin{aligned}
\tilde{g}(t)\tilde{h}(t) &= \left(\sum_{T' \in \bar{\mathbb{T}}^W} t^{|T'|} \mathcal{V}_{T'} \right) \left(\sum_{T'' \in \bar{\mathbb{H}}^W} \beta(T'') t^{|T''|-1} \mathcal{V}_{T''} \right) \\
&= \sum_{T \in \bar{\mathbb{T}}^W} t^{|T|-1} \left(\sum_{\substack{C \in \mathcal{C}(T) \\ R_C(T) \in \bar{\mathbb{H}}^W}} \beta(R_C(T)) \right) \mathcal{V}_T.
\end{aligned}$$

Note that, the set of all admissible cuts C such that $R_C(T) \in \bar{\mathbb{H}}^W$ is in 1-1 correspondence with the set of rooted subtrees S of T with $S \in \bar{\mathbb{H}}^W$. But any subtree S of T with $S \in \bar{\mathbb{H}}^W$ is completely determined by the unique leaf of S . Therefore, the set of all admissible cuts C such that $R_C(T) \in \bar{\mathbb{H}}^W$ is in 1-1 correspondence with the set of non-root vertices of T . With this observation, we have, for any $T \in \bar{\mathbb{T}}^W$,

$$(4.20) \quad \sum_{\substack{C \in \mathcal{C}(T) \\ R_C(T) \in \bar{\mathbb{H}}^W}} \beta(R_C(T)) = |T|.$$

Hence, combining the two equations above, we get

$$\tilde{g}(t)\tilde{h}(t) = \sum_{T \in \bar{\mathbb{T}}^W} t^{|T|-1} |T| \mathcal{V}_T = \frac{d\tilde{g}(t)}{dt},$$

which is Eq. (2.5) for the system $\Omega_{\mathbb{T}}^W$. $\square$

Now, by the universal property of the NCS system $(\mathcal{NSym}, \Pi)$ from NCSFs in Theorem 2.6, we have the following correspondence between NCSFs and W -labeled rooted trees.

Theorem 4.6. *For any nonempty $W \subseteq \mathbb{N}^+$, there exists a unique homomorphism $\mathcal{T}_W : \mathcal{NSym} \rightarrow \mathcal{H}_{GL}^W$ of graded K -Hopf algebras such that $\mathcal{T}_W^{\times 5}(\Pi) = \Omega_{\mathbb{T}}^W$.*

In particular, we have the following correspondence from the NCSFs in Π to the elements in $\Omega_{\mathbb{T}}^W$:

$$(4.21) \quad \mathcal{T}_W(\Lambda_m) = \sum_{T \in \mathbb{S}_m^W} (-1)^{o(T)+|T|} \mathcal{V}_T,$$

$$(4.22) \quad \mathcal{T}_W(S_m) = \sum_{T \in \mathbb{T}_m^W} \mathcal{V}_T,$$

$$(4.23) \quad \mathcal{T}_W(\Psi_m) = \sum_{T \in \mathbb{H}_m^W} \beta_T \mathcal{V}_T,$$

$$(4.24) \quad \mathcal{T}_W(\Phi_m) = m \sum_{T \in \mathbb{P}_m} \theta_T \mathcal{V}_T,$$

$$(4.25) \quad \mathcal{T}_W(\Xi_m) = \sum_{T \in \mathbb{P}_m} \gamma_T \mathcal{V}_T,$$

for any $m \geq 1$.

Proof: Note that the coefficients of t^m ($m \geq 1$) of the generating function $\tilde{h}(t)$ (see Eq. (4.3)) are all primitive elements of the Hopf algebra $\mathcal{H}_{GL}^W$, since they are linear combinations of chains. Then, by Theorem 2.6, (b), we have a unique homomorphism $\mathcal{T}_W : \mathcal{NSym} \rightarrow \mathcal{H}_{GL}^W$ of K -Hopf algebras such that $\mathcal{T}_W^{\times 5}(\Pi) = \Omega_{\mathbb{T}}^W$. In particular, we have $\mathcal{T}_W(\lambda(t)) = \tilde{f}(t)$ which is same as Eq. (4.21) for any $m \geq 1$. Note that, both sides of Eq. (4.21) have weight m in $\mathcal{NSym}$ and $\mathcal{H}_{GL}^W$, respectively. Also note that the gradings of $\mathcal{NSym}$ and $\mathcal{H}_{GL}^W$ are given by the weights of NCSFs and W -labeled rooted trees, respectively, and $\mathcal{NSym}$ is the free algebra generated by Λ_m ($m \geq 1$). By the facts above, it is easy to check that $\mathcal{T}_W$ also preserves the gradings. $\square$

Note that the graded duals of $\mathcal{NSym}$ and $\mathcal{H}_{GL}^W$ are the graded K -Hopf algebras $\mathcal{QSym}$ of quasi-symmetric functions and the Connes-Kreimer Hopf algebra $\mathcal{H}_{CK}^W$, respectively. Since the K -Hopf algebra homomorphism $\mathcal{T}_W : \mathcal{NSym} \rightarrow \mathcal{H}_{GL}^W$ preserves the gradings, we can take the graded duals and get the following correspondence.

Corollary 4.7. *For any non-empty $W \subseteq \mathbb{N}^+$, $\mathcal{T}_W^* : \mathcal{H}_{CK}^W \rightarrow \mathcal{QSym}$ is a homomorphism of graded K -Hopf algebras.*

Finally, let us end this section with the following two remarks.

Remark 4.8. *As we mentioned earlier in Remark 2.3, by applying the specialization $\mathcal{T}_W : \text{NSym} \rightarrow \mathcal{H}_{GL}^W$ in Theorem 4.6, we will get a host of identities for the W -rooted trees on the right hand sides of Eqs. (4.21)–(4.25) from the identities of the NCSFs on the left hands. We believe some of these identities are also interesting from the aspect of combinatorics of rooted trees. For example, it is not obvious at all that the invariants of rooted trees given by coefficients of the generating functions $\tilde{f}(t)$, $\tilde{d}(t)$, $\tilde{h}(t)$ and $\tilde{m}(t)$ can be obtained by evaluating the corresponding (noncommutative) polynomials of NCSFs at coefficients of $\tilde{g}(t)$ which is just the trivial invariant of rooted trees whose value at any rooted tree is always 1.*

Note that the same problem has been studied in detail in [Z4] for the differential operators in the NCS over the differential operator algebras constructed in [Z3]. But, in order to keep this paper in a certain size, we will skip the discussions on these identities and refer the interested reader to [Z4] for similar discussions.

Remark 4.9. *In the followed paper [Z5], by using some relations of the NCS system $(\mathcal{H}_{GL}^W, \Omega_{\mathbb{T}}^W)$ with the NCS systems constructed in [Z3] over differential operator algebras, it will be shown that, when $W = \mathbb{N}^+$, the K -Hopf algebra homomorphism $\mathcal{T}_W : \text{NSym} \rightarrow \mathcal{H}_{GL}^W$ in Theorem 4.6 is actually an embedding. So, in this case, $\mathcal{T}_W^* : \mathcal{H}_{CK}^W \rightarrow \mathcal{QSym}$ in Corollary 4.7 is a surjective graded K -Hopf algebra homomorphism.*

It seems that, the K -Hopf algebra homomorphism $\mathcal{T}_W : \text{NSym} \rightarrow \mathcal{H}_{GL}^W$ is injective for any non-empty label set W . But we will leave this for future investigations.

5. A Combinatorial Interpretation of the Constants θ_T

In this section, we give a combinatorial interpretation for the constants θ_T ($T \in \mathbb{T}$) in Definition 4.2 which have been used in the construction of $\tilde{d}(t)$ for the NCS system $\Omega_{\mathbb{T}}^W$ over $\mathcal{H}_{GL}^W$. We first recall the strict order polynomials and the order polynomials of finite posets (partially ordered sets) (see [St1]). Then, by applying some of results proved in [WZ] for the strict order polynomials of rooted forests and the well-known Reciprocity Relation (see Proposition 5.1) between the strict order polynomials and the order polynomials of finite posets, we show in Proposition 5.8 that, for any $T \in \bar{\mathbb{T}}$, θ_T is same as the coefficient of s of the order polynomial of the rooted forest $B_-(T)$.

First, recall that, a *poset* is a set P with a partial order defined for its elements. Note that, for any unlabeled rooted forest $F \in \mathbb{F}$, the set

of the vertices of F has a natural partial order, namely, $u \leq v$ if $u = v$ or u and v are connected by a path with v being further away for the root of the connected component. With this partial order, the set of the vertices of F forms a finite poset, which we will still denote by F .

For any $n \geq 1$, let $[n]$ denote the totally ordered set $\{1, 2, \dots, n\}$. For any finite poset P , a map $f : P \rightarrow [n]$ is said to be *strictly order-preserving* (resp. *order-preserving*) if, for any $a, b \in P$ with $a < b$ in P , then $f(a) < f(b)$ (resp. $f(a) \leq f(b)$). It is well-known that, for each finite poset P , there exists a unique polynomial $\bar{\Omega}(P, s)$ (resp. $\Omega(P, s)$) in formal variable s such that, for any $n \geq 1$, $\bar{\Omega}(P, n)$ (resp. $\Omega(P, n)$) equals to the number of strict order-preserving (resp. *order-preserving*) from P to $[n]$. The strict order polynomial and the order polynomial of finite posets are related by the so-called reciprocity relation.

Proposition 5.1. (Reciprocity Relation) *For any fixed finite poset, we have*

$$(5.1) \quad \Omega(P, s) = (-1)^{|P|} \bar{\Omega}(P, -s),$$

where $|P|$ denotes the cardinal number of the finite set P .

For a proof of this remarkable result, see Corollary 4.5.15 in [St1].

By Corollary 3.15 in [WZ] (also see Example 4.4 there) and Eq. (5.1) above, we get the (strict) order polynomials of chains and shrubs as follows.

Example 5.2. *For any $m \geq 1$, let C_m be the chain of height $m - 1$ and S_m the shrub with m leaves. Then we have*

$$(5.2) \quad \bar{\Omega}(C_m, s) = \binom{s}{m} = \frac{s(s-1) \cdots (s-m+1)}{m!},$$

$$(5.3) \quad \Omega(C_m, s) = (-1)^m \binom{-s}{m} = \binom{s+m-1}{m},$$

$$(5.4) \quad \bar{\Omega}(S_m, s) = \int_0^s B_m(u) du, = \frac{B_{m+1}(s) - B_{m+1}(0)}{m+1},$$

$$(5.5) \quad \Omega(S_m, s) = (-1)^{m+1} \int_0^{-s} B_m(u) du,$$

where $B_m(u)$ ($m \geq 1$) are the Bernoulli polynomials which are defined by the generating function

$$\frac{xe^{ux}}{e^x - 1} = 1 + \sum_{m=1}^{\infty} B_m(u) \frac{x^m}{m!}.$$

Proposition 5.3. (a) *For any finite poset P , $\bar{\Omega}(P, 0) = 0$.*

(b) For any rooted forest $F = T_1 T_2 \cdots T_m$ with $T_i \in \mathbb{T}$ ($1 \leq i \leq m$), we have

$$(5.6) \quad \bar{\Omega}(F, s) = \bar{\Omega}(T_1, s) \bar{\Omega}(T_2, s) \cdots \bar{\Omega}(T_m, s).$$

(c) For any unlabeled rooted tree T with $T = B_+(F)$, we have

$$(5.7) \quad \Delta \bar{\Omega}(T, s) = \bar{\Omega}(F, s),$$

$$(5.8) \quad \nabla \Omega(T, s) = \Omega(F, s),$$

where Δ and ∇ are the linear operators from $K[s] \rightarrow K[s]$ defined by, for any $f(s) \in K[s]$, $\Delta f(s) = f(s+1) - f(s)$ and $\nabla f(s) = f(s) - f(s-1)$, respectively.

(a) is well-known, for example, it can be easily proved by the recurrent formulas in [SWZ] for the (strict) order polynomials. (b) follows directly from the definition of the strict order polynomials. (c) was first proved by J. Shareshian (unpublished). For a proof of Eq. (5.7), see Theorem 4.5 in [WZ]. Eq. (5.8) can be proved similarly as Eq. (5.7). For more studies on these properties of the (strict) order polynomials, see [Z1] and [SWZ].

Now, for any finite poset P , we define

$$(5.9) \quad \phi_P := \left. \frac{d}{ds} \bar{\Omega}(P, s) \right|_{s=0},$$

$$(5.10) \quad \varphi_P := \left. \frac{d}{ds} \Omega(P, s) \right|_{s=0}.$$

By Proposition 5.1 and 5.3, it is easy to see that we have the following corollary.

Corollary 5.4. (a) For any finite poset P , we have $\phi_P = (-1)^{|P|-1} \varphi_P$.

(b) For any rooted forest $F \in \mathbb{F}$ with more than one connected component, we have $\phi_F = \varphi_F = 0$.

The following proposition have been proved in [WZ]. But note that the definition of $T_{\vec{e},k}$ we adapt here is different form the one used in [WZ]. So the equations in the proposition below have been modified accordingly.

Proposition 5.5. (a) The constants $\{\phi_T \mid T \in \mathbb{T}\}$ satisfy, and are uniquely determined by

$$\phi_{T=\circ} = 1,$$

$$\phi_T = - \sum_{k=2}^{v(T)} \frac{1}{k!} \sum_{\substack{\vec{e}=(e_1, \dots, e_{k-1}) \in E(T)^{k-1} \\ e_1 \succ \dots \succ e_{k-1}}} \phi_{B_-(T_{\vec{e},1})} \cdots \phi_{B_-(T_{\vec{e},k-1})} \phi_{T_{\vec{e},k}},$$

when $v(T) \geq 2$.

(b) For any $T \in \mathbb{T}$, we have

$$\bar{\Omega}(T, s) = \phi_T s + \sum_{k=2}^{v(T)} \frac{s^k}{k!} \sum_{\substack{\vec{e}=(e_1, \dots, e_{k-1}) \in E(T)^{k-1} \\ e_1 \succ \dots \succ e_{k-1}}} \phi_{B_-(T_{\vec{e},1})} \cdots \phi_{B_-(T_{\vec{e},k-1})} \phi_{T_{\vec{e},k}}.$$

By Proposition 5.5, Corollary 5.4, (a) and Proposition 5.1, it is easy to check that the order polynomials $\Omega(T, s)$ of rooted trees T can be obtained as follows.

Corollary 5.6. (a) The constants $\{\varphi_T | T \in \mathbb{T}\}$ satisfy and are uniquely determined by

$$\begin{aligned} \varphi_{T=o} &= 1, \\ \varphi_T &= \sum_{k=2}^{v(T)} \frac{(-1)^k}{k!} \sum_{\substack{\vec{e}=(e_1, \dots, e_{k-1}) \in E(T)^{k-1} \\ e_1 \succ \dots \succ e_{k-1}}} \varphi_{B_-(T_{\vec{e},1})} \cdots \varphi_{B_-(T_{\vec{e},k-1})} \varphi_{T_{\vec{e},k}} \end{aligned}$$

when $v(T) \geq 2$.

(b) For any $T \in \mathbb{T}$, we have

(5.11)

$$\Omega(T, s) = \varphi_T s + \sum_{k=2}^{v(T)} \frac{s^k}{k!} \sum_{\substack{\vec{e}=(e_1, \dots, e_{k-1}) \in E(T)^{k-1} \\ e_1 \succ \dots \succ e_{k-1}}} \varphi_{B_-(T_{\vec{e},1})} \cdots \varphi_{B_-(T_{\vec{e},k-1})} \varphi_{T_{\vec{e},k}}.$$

Note that, from the definition of the order polynomials, we have $\Omega(P, 1) = 1$ for any finite poset P . Using this fact and evaluating $\Omega(T, s)$ in Eq. (5.11) at $s = 1$, we get another recurrent formula for the constants φ_T of rooted trees.

Corollary 5.7. The constants $\{\varphi_T | T \in \mathbb{T}\}$ satisfy and are uniquely determined by

$$\begin{aligned} \varphi_{T=o} &= 1, \\ (5.12) \end{aligned}$$

$$\varphi_T = 1 - \sum_{k=2}^{v(T)} \frac{1}{k!} \sum_{\substack{\vec{e}=(e_1, \dots, e_{k-1}) \in E(T)^{k-1} \\ e_1 \succ \dots \succ e_{k-1}}} \varphi_{B_-(T_{\vec{e},1})} \cdots \varphi_{B_-(T_{\vec{e},k-1})} \varphi_{T_{\vec{e},k}}.$$

Now, we consider the constants θ_T ($T \in \mathbb{T}$) defined in Definition 4.2 in Section 4 and prove the following main result of this section.

Proposition 5.8. *For any $T \in \mathbb{T}$ with $T = B_+(F)$, we have*

$$(5.13) \quad \theta_T = \varphi_F.$$

Proof: For any $T \in \mathbb{T}$ with $T = B_+(F)$, we set $\tilde{\theta}_T := \varphi_F$. For convenience, we also set $\tilde{\theta}_\circ = 0$. We need show that $\tilde{\theta}_T = \theta_T$ for any $T \in \mathbb{T}$.

Note first that, if T the singleton $\circ$ or a non-primitive rooted tree, i.e. F is the empty or has at least two connected components, by the definition of θ_T in Definition 4.2 and Corollary 5.4, (b), we have $\theta_T = \tilde{\theta}_T = 0$. To show that $\tilde{\theta}_T = \theta_T$ for all primitive rooted trees T , it will be enough to show that, $\tilde{\theta}_T$ ($T \in \mathbb{P}$) also satisfies the recurrent relations in Definition 4.2.

We use the mathematical induction on $v(T)$. First, for the case $v(T)$, i.e. $T = B_+(\circ) = C_2$. Since $\Omega(\circ, s) = s$, $\tilde{\theta}_T = \varphi_\circ = 1$. While θ_T is defined to be 1 in Definition 4.2. Hence, $\theta_T = \tilde{\theta}_T$ in this case.

Now, assume $T = B_+(T')$ with $T' \in \mathbb{T}$ and $v(T') \geq 2$. Applying Eq. (5.12) to T' , we have

$$\varphi_{T'} = 1 - \sum_{k=2}^{v(T')} \frac{1}{k!} \sum_{\substack{\vec{e}=(e_1, \dots, e_{k-1}) \in E(T')^{k-1} \\ e_1 \succ \dots \succ e_{k-1}}} \varphi_{B_-(T_{\vec{e},1})} \cdots \varphi_{B_-(T_{\vec{e},k-1})} \varphi_{T_{\vec{e},k}}.$$

Note that, we can identify the set of $\vec{e} = (e_1, \dots, e_{k-1}) \in E(T')^{k-1}$ with the set of $\vec{e} = (e_1, \dots, e_{k-1}) \in E(T)^{k-1}$ such that $T_{\vec{e},k} \neq \circ$. With these observations, by replacing the constants φ 's by $\tilde{\theta}$'s in the summation of the equation above, we have

$$\tilde{\theta}_T = 1 - \sum_{k=2}^{v(T)} \frac{1}{k!} \sum_{\substack{\vec{e}=(e_1, \dots, e_{k-1}) \in E(T)^{k-1} \\ e_1 \succ \dots \succ e_{k-1} \\ T_{\vec{e},k} \neq \circ}} \tilde{\theta}_{T_{\vec{e},1}} \tilde{\theta}_{T_{\vec{e},2}} \cdots \tilde{\theta}_{T_{\vec{e},k}}$$

Applying the induction assumption to $T_{\vec{e},j}$'s and using the fact that $\tilde{\theta}_\circ = 0$:

$$= 1 - \sum_{k=2}^{v(T)} \frac{1}{k!} \sum_{\substack{\vec{e}=(e_1, \dots, e_{k-1}) \in E(T)^{k-1} \\ e_1 \succ \dots \succ e_{k-1}}} \tilde{\theta}_{T_{\vec{e},1}} \tilde{\theta}_{T_{\vec{e},2}} \cdots \tilde{\theta}_{T_{\vec{e},k}}.$$

Therefore, the constants $\{\tilde{\theta}_T \mid T \in \mathbb{P}\}$ also satisfy the recurrent relations of $\{\theta_T \mid T \in \mathbb{P}\}$ in Definition 4.2. Hence, we have $\theta_T = \tilde{\theta}_T = \varphi_T$ for any $T \in \mathbb{T}$. $\square$

For an interpretation of the constant ϕ_T , which we have shown is same as $(-1)^{v(T)-1}\varphi_T = (-1)^{v(T)-1}\theta_{B_+(T)}$, in terms of the numbers of chains with fixed lengths in the lattice of the ideals of the poset T , see Lemma 2.8 in [SWZ].

Corollary 5.9. *For any $T \in \mathbb{P}$, we have*

(5.14)

$$\nabla\Omega(T, s) = \theta_T s + \sum_{k=2}^{v(T)} \frac{s^k}{k!} \sum_{\substack{\vec{e}=(e_1, \dots, e_{k-1}) \in E(T)^{k-1} \\ e_1 \succ \dots \succ e_{k-1}}} \theta_{B_-(T_{\vec{e},1})} \cdots \theta_{B_-(T_{\vec{e},k-1})} \theta_{T_{\vec{e},k}}.$$

In particular, θ_T is also the coefficient of s of the polynomial $\nabla\Omega(T, s)$.

Proof: First, we write $T = B_+(T')$ with $T' \in \mathbb{T}$. By Eq. (5.8) and Proposition 5.5, (b), we have

$$\begin{aligned} \nabla\Omega(T, s) &= \Omega(T', s) \\ &= \varphi_{T'} s + \sum_{k=2}^{v(T')} \frac{s^k}{k!} \sum_{\substack{\vec{e}=(e_1, \dots, e_{k-1}) \in E(T')^{k-1} \\ e_1 \succ \dots \succ e_{k-1}}} \varphi_{B_-(T_{\vec{e},1})} \cdots \varphi_{B_-(T_{\vec{e},k-1})} \varphi_{T'_{\vec{e},k}}. \end{aligned}$$

Then, applying Eq. (5.13) and replacing the constants φ 's in the sum above by the constant θ 's, we get Eq. (5.14). $\square$

REFERENCES

- [A] E. Abe, *Hopf Algebras*. Translated from the Japanese by Hisae Kinoshita and Hiroko Tanaka. Cambridge Tracts in Mathematics, 74. Cambridge University Press, Cambridge-New York, 1980. [MR0594432].
- [ABS] M. Aguiar, N. Bergeron and F. Sottile, *Combinatorial Hopf Algebras and Generalized Dehn-Sommerville Relations*, Compos. Math. 142 (2006), no. 1, 1–30. [MR2196760]. See also math.CO/0310016.
- [BCW] H. Bass, E. Connell, D. Wright, *The Jacobian conjecture, reduction of degree and formal expansion of the inverse*. Bull. Amer. Math. Soc. **7**, (1982), 287–330. [MR 83k:14028].
- [CK] A. Connes and D. Kreimer, *Hopf Algebras, Renormalization and Non-commutative Geometry*, Comm. Math. Phys. 199 (1998), no. 1, 203–242. [MR1660199]. See also hep-th/9808042.
- [DFT] G. Duchamp; A. Klyachko; H. Hivert and J.-Y Thibon, *Noncommutative symmetric functions. VI. Free quasi-symmetric functions and related algebras*, Internat. J. Algebra Comput. 12 (2002), no. 5, 671–717. [MR1935570].

- [DKKT] G. Duchamp; A. Klyachko; D. Krob and J.-Y Thibon, *Noncommutative symmetric functions. III. Deformations of Cauchy and convolution algebras*, Lie computations (Marseille, 1994). Discrete Math. Theor. Comput. Sci. 1 (1997), no. 1, 159–216. [MR1605038].
- [E] A. van den Essen, *Polynomial automorphisms and the Jacobian conjecture*. Progress in Mathematics, 190. Birkhäuser Verlag, Basel, 2000. [MR1790619].
- [F] L. Foissy, *Les algèbres de Hopf des arbres enracinés décorés I, II*, Bull. Sci. Math. 126 (2002), no. 3, 193–239 & no. 4, 249–288. [MR1909461] & [MR1909461]. See also math.QA/0105212.
- [GKLLRT] I. M. Gelfand; D. Krob; A. Lascoux; B. Leclerc; V. S. Retakh and J.-Y. Thibon, *Noncommutative symmetric functions*. Adv. Math. 112 (1995), no. 2, 218–348. [MR1327096]. See also hep-th/9407124.
- [Ge] I. Gessel, *Multipartite P -partitions and inner products of skew Schur functions*, Comtemp. Math. **34** (1984), 289–301. [MR0777705].
- [GL] R. Grossman and R. G. Larson, *Hopf-Algebraic structure of families of Trees*, J. Algebra **126** (1989), no.1, 184–210. [MR1023294].
- [H] M. E. Hoffman, *Combinatorics of Rooted Trees and Hopf Algebras*, Trans. Amer. Math. Soc. **355** (2003), no. 9, 3795–3811. [MR1990174]. See also math.CO/0201253.
- [Knu] D. Knutson, *λ -Rings and the Representation Theory of the Symmetric Groups*, Lecture Notes in Mathematics, Vol. **308**, Springer-Verlag, Berlin-New York, 1973. [MR0364425].
- [Kr] D. Kreimer, *On the Hopf Algebra Structure of Perturbative Quantum Field Theories*, Adv. Theor. Math. Phys. **2.2** (1998) 303–334. [MR1633004]. See also math.QA/9707029.
- [KLT] D. Krob; B. Leclerc and J.-Y Thibon, *Noncommutative symmetric functions. II. Transformations of alphabets*, Internat. J. Algebra Comput. 7 (1997), no. 2, 181–264. [MR1433196].
- [KT1] D. Krob and J.-Y Thibon, *Noncommutative symmetric functions. IV. Quantum linear groups and Hecke algebras at $q = 0$* , J. Algebraic Combin. 6 (1997), no. 4, 339–376. [MR1471894].
- [KT2] D. Krob and J.-Y Thibon, *Noncommutative symmetric functions. V. A degenerate version of $U_q(\mathfrak{gl}_N)$* , Internat. J. Algebra Comput. 9 (1999), no. 3-4, 405–430. [MR1723475].
- [Mc] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, Second edition. With contributions by A. Zelevinsky. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1995. [MR1354144].
- [MR] C. Malvenuto and C. Reutenauer, *Duality between Quasi-Symmetric Functions and the Solomon Descent Algebra*, J. Algebra **177** (1995), no.3, 967–982. [MR1358493].
- [Mo] S. Montgomery, *Hopf Algebras and Their Actions on Rings*, CBMS Regional Conference Series in Mathematics, **82**, Providence, R. I. 1993. [MR1243637].
- [SWZ] J. Shareshian, D. Wright and W. Zhao, *A New Approach to Order Polynomials of Labeled Posets and Their Generalizations*, math.CO/0311426.

- [St1] Richard P. Stanley, *Enumerative Combinatorics I*, Cambridge University Press, 1997. [MR 98a:05001].
- [St2] Richard P. Stanley, *Enumerative Combinatorics II*, Cambridge University Press, 1999. [MR 2000k:05026].
- [T] J.-Y Thibon, *Lectures on noncommutative symmetric functions*, Interaction of combinatorics and representation theory, 39–94, MSJ Mem., 11, Math. Soc. Japan, Tokyo, 2001. [MR1862149].
- [WZ] D. Wright and W. Zhao, *D-log and formal flow for analytic isomorphisms of n -space*. Trans. Amer. Math. Soc. 355 (2003), no. 8, 3117–3141. [MR1974678]. See also math.CV/0209274.
- [Z1] W. Zhao, *A family of invariants of rooted forests*, J. Pure Appl. Algebra **186** (2004), no. 3, 311–327. [MR 2025592]. See also math.CO/0211095.
- [Z2] W. Zhao, *Noncommutative Symmetric Systems over Associative Algebras*, J. Pure Appl. Algebra, **210**, no. 2, 363–382. See also math.CO/0509133.
- [Z3] W. Zhao, *Differential Operator Specializations of Noncommutative Symmetric Functions*, Advances in Mathematics, **214**, no. 2, 639–665. See also math.CO/0509134.
- [Z4] W. Zhao, *Noncommutative Symmetric Functions and the Inversion Problem*, math.CV/0509135. Submitted.
- [Z5] W. Zhao, *NCS Systems over Differential Operator Algebras and the Grossman-Larson Hopf Algebra of Labeled Rooted Trees*, preprint, math.CO/0509138. Submitted.

DEPARTMENT OF MATHEMATICS, ILLINOIS STATE UNIVERSITY, NORMAL,
IL 61790-4520.

E-mail: wzhao@ilstu.edu.